

*Trichoptera, &c., in North Devon.*—In Ent. Mo. Mag., vol. xxvi, p. 316, Mr. McLachlan records the species of *Trichoptera* observed by himself and two friends on Exmoor in September, 1890, and by Mr. Stanley Edwards near Lynton, North Devon, from August 8th to September 18th, 1890. During a short stay at Lynton from June 9th to 22nd, although most of my time was otherwise engaged, I was able to add the following species to the Lynton list, and to assure myself of a rich harvest next year:—*Lasiocephala basalis*, *Dipterotreron felix*, *Psychomyia pusilla*, *Lype phæopa*, and *Glossosoma Boltoni*, very large and dark. Of the species previously recorded, as above, *Hydropsyche instabilis* was out in countless myriads, and *Philopotamus montanus* was generally abundant and very brightly coloured. I also took two *Osmylus chrysops*. In the East Lynn I secured a short series of *Perla maxima* and a few *Perla cephalotes*, the latter species not being detected until Mr. K. J. Morton kindly drew my attention to the differences between the species. *Chloroperla grammatica* and *Isopteryx tripunctata* were common, as also was a *Leuctra* not yet quite determined. I trust next year to largely increase the list of Lynton *Neuroptera*.—C. A. BRIGGS, 55, Lincoln's Inn Fields: September 2nd, 1896.

*Notes on a few Neuroptera from the Tyrol.*—The greater part of last month and the commencement of the present were spent by my friends Messrs. Champion and Lemann in the Tyrol. They kindly collected a few *Neuroptera* (in the broad sense) which they handed over to me, and as an item towards a local fauna that may possibly be of use hereafter, I have given a list of them.

The approximate elevations of the localities mentioned are as follows:—Cortina, 3900 ft.; Mendel, 4400 feet; Campiglio, 4900 ft.; Sorapis, 6-7000 ft.

TRICHOPTERA.—*Limnophilus rhombicus*, L., *L. flavicornis*, F. (the anterior wings strongly sprinkled with dark markings), and *L. ignavus*, McL. (one ♀); *Sericostoma* sp. (♀ only); *Odontocerum albicorne*, Sc.; *Philopotamus ludificatus*, McL.; *Tinodes dives*, Pict. All from Mendel.

PLANIPENNIA.—*Panorpa alpina*, Rbr., Sorapis and Mendel, *P. communis*, L., Campiglio and Mendel, *P. germanica*, L., var. *apicalis*, Steph., Cortina, *P. cognata*, Rbr., Campiglio; *Rhaphidia flavipes*, Stein, Campiglio; *Osmylus chrysops*, L., Mendel (common); *Hemerobius nitidulus*, F., Mendel, *H. concinnus*, Steph., var. *4-fasciatus*, Reut., Campiglio and Mendel; *Megalomus hirtus*, L., Campiglio; *Chrysopa ventralis*, C., *C. prasina*, Burm. (*aspersa*, Wesm.), and *C. vulgaris*, Schnd., Mendel.

PSEUDO-NEUROPTERA.—*Perla maxima*, Sc., Campiglio (three ♂, in colour nearly as dark as Scotch examples, but with ample wings); *Nemoura variegata*, Oliv., Campiglio; *Sympetrum striolatum*, Charp., and *S. meridionale*, Selys (one), Campiglio; *Somatochlora arctica*, Zett., Mendel (a pair, ♂ ♀); *Cordulegaster bidentatus*, Selys, Campiglio (one ♀).—R. McLACHLAN, Lewisham, London: August 19th, 1896.

*Coleoptera at Broadstairs.*—While staying with friends near Broadstairs at the beginning of August I discovered that some of their early potatoes still remained in the ground, so I promptly relieved the gardener of the task of lifting them, and was delighted to take from the decaying seed almost all the *Coleoptera* which used to occur to me, under similar circumstances, in the same neighbourhood ten years